

Light Like Segment Compactification and Braneworlds with Dynamical String Tension

E.I. Guendelman

Department of Physics, Ben-Gurion University of the Negev, Beer-Sheva, Israel

Frankfurt Institute for Advanced Studies, Giersch Science Center, Campus Riedberg,
Frankfurt am Main, Germany

Bahamas Advanced Studies Institute and Conferences, 4A Ocean Heights, Hill View
Circle, Stella Maris, Long Island, The Bahamas

E-mail: guendel@bgu.ac.il,

Abstract

There is great interest in the construction of brane worlds, where matter and gravity are forced to be effective only in a lower dimensional surface, the "brane". How these could appear as a consequence of string theory is a crucial question and this has been widely discussed. Here we will examine a distinct scenario that appears in dynamical string tension theories and where string tension is positive between two surfaces separated by a short distance and at the two surfaces themselves the string tensions become infinite, therefore producing an effective confinement of the strings and therefore of all matter and gravity to the space between these two surfaces, which is in fact a new type of stringy brane world scenario. The specific model studied is in the context of the modified measure formulation of the string where tension appears as an additional dynamical degree of freedom and these tensions are not universal, but rather each string generates its own tension, which can have a different value for each string. We consider a new background field that can couple to these strings, the "tension scalar" is capable then of changing locally along the world sheet and then the value of the tension of the extended object changes accordingly. When many types of strings probing the same region of space are considered this tension scalar is constrained by the requirement of quantum conformal invariance. For the case of two types of strings probing the same region of space with different dynamically generated tensions, there are two different metrics, associated to the different strings, that have to satisfy vacuum Einstein's equations and the consistency of these two Einstein's equations determine the tension scalar. The universal metric, common to both strings generically does not satisfy Einstein's equation. The two metrics considered here are flat space in Minkowski space and flat space after a special conformal transformation and the tension field behaves in such a way that strings are confined inside a light like segment or alternatively as expanding braneworlds where the strings are confined between two expanding bubbles separated by a very small distance at large times.

1 Introduction

The basic idea of the brane worlds is that the universe is restricted to a brane inside a higher-dimensional space, called the "bulk" . In this model, at least some of the extra dimensions are extensive (possibly infinite), and other branes may be moving through this bulk. Some of the first braneworld models were developed by Rubakov and Shaposhnikov [1], Visser [2], Randall and Sundrum [3], [4], Pavsic [5], Gogberashvili [6]. At least some of these models are motivated by string theory. For discussions and a review of braneworlds in string theory see for example [7], our approach will be very different to the present standard approaches to braneworlds in the context of string theories however. In our approach a dynamical string tension is required. Our scenario could be enriched by incorporating aspects of the more traditional braneworlds, but these aspects will be ignored here to simplify the discussion.

String Theories have been considered by many physicists for some time as the leading candidate for the theory everything, including gravity, the explanation of all the known particles that we know and all of their known interactions (and probably more) [8]. According to some, one unpleasant feature of string theory as usually formulated is that it has a dimension full parameter, in fact, its fundamental parameter , which is the tension of the string. This is when formulated the most familiar way. The consideration of the string tension as a dynamical variable, using the modified measures formalism, which was previously used for a certain class of modified gravity theories under the name of Two Measures Theories or Non Riemannian Measures Theories, see for example [9, 10, 12, 11, 13, 14, 15, 16]. In the context of this paper, it is also interesting to mention that the modified measure approach has also been used to construct braneworld scenarios [17]

When applying these principles to string theory, this leads to the modified measure approach to string theory, where rather than to put the string tension by hand it appears dynamically.

This approach has been studied in various previous works [18, 19, 20, 21, 22, 23, 24, 25]. See also the treatment by Townsend and collaborators for dynamical string tension [27, 28].

In our most recent papers on the subject [25], we have also introduced the "tension scalar", which is an additional background fields that can be introduced into the theory for the bosonic case (and expected to be well defined for all types of superstrings as well) that changes the value of the tension of the extended object along its world sheet, we call this the tension scalar for obvious reasons. Before studying issues that are very special of this paper we review some of the material contained in previous papers, first present the string theory with a modified measure and containing also gauge fields that like in the world sheet, the integration of the equation of motion of these gauge fields gives rise to a dynamically generated string tension, this string tension may differ from one string to the other.

Then we consider the coupling of gauge fields in the string world sheet to currents in this world sheet, as a consequence this coupling induces variations of the tension along the world sheet of the string. Then we consider a bulk scalar and how this scalar naturally can induce this world sheet current that couples to the internal gauge fields. The integration of the equation of motion of the internal gauge field lead to the remarkably simple equation that the local value of the tension along the string is given by $T = e\phi + T_i$, where e is a coupling constant that defines the coupling of the bulk

scalar to the world sheet gauge fields and T_i is an integration constant which can be different for each string in the universe.

Then each string is considered as an independent system that can be quantized. We take into account the string generation by introducing the tension as a function of the scalar field as a factor inside a Polyakov type action with such string tension, then the metric and the factor $g\phi + T_i$ enter together in this effective action, so if there was just one string the factor could be incorporated into the metric and the condition of world sheet conformal invariance will not say very much about the scalar ϕ , but if many strings are probing the same regions of space time, then considering a background metric $g_{\mu\nu}$, for each string the "string dependent metric" $(\phi + T_i)g_{\mu\nu}$ appears and in the absence of other background fields, like dilaton and antisymmetric tensor fields, Einstein's equations apply for each of the metrics $(\phi + T_i)g_{\mu\nu}$, considering two types of strings with different tensions. We call $g_{\mu\nu}$ the universal metric, which in fact does not necessarily satisfy Einstein's equations.

In the case of the flat space for the string associated metrics, in the Milne representation, for the case of two types of string tensions, we study the case where the two types of strings have positive string tensions, as opposed to our previous work [25] where we found solutions with both positive and negative string tensions. At the early universe the negative string tension strings tensions are large in magnitude, but approach zero in the late universe and the positive string tensions appear for the late universe with their tension approaching a constant value at the late universe. These solutions are absolutely singularity free.

In contrast, we have studied also [26] the case of very different solutions and where both type of strings have positive tensions, then these are singular, they cannot be continued before a certain time (that corresponded to a bounce in our previous work [25]). Here, at the origin of time, the string tensions of both types of strings approach plus infinity, so this opens the possibility of having no Hagedorn temperature in the early universe and latter on in the history of the universe as well for this type of string cosmology scenario. Also the tensions can become infinity at a certain location in the warped coordinate in a warped scenario.

Here we will study a situation where we consider the metrics $(\phi + T_i)g_{\mu\nu}$, considering two types of strings tensions. The two metrics will again satisfy Einstein's equations and the two metrics will represent Minkowski space and Minkowski space after a special conformal transformation. In this case, the location where the two types of strings acquire an infinite tension is given by two surfaces. If the vector that defines the special conformal transformation is light like, these two surface are plane, parallel to each other and both move with the speed of light. If the vector is time like of space like the two surfaces are spherical and expanding and the distance between them approaches zero at large times (positive or negative). In both cases this represents a genuine brane world scenario.

2 The Modified Measure Theory String Theory

The standard world sheet string sigma-model action using a world sheet metric is [29], [30], [31]

$$S_{sigma-model} = -T \int d^2\sigma \frac{1}{2} \sqrt{-\gamma} \gamma^{ab} \partial_a X^\mu \partial_b X^\nu g_{\mu\nu}. \quad (1)$$

Here γ^{ab} is the intrinsic Riemannian metric on the 2-dimensional string worldsheet and $\gamma = \det(\gamma_{ab})$; $g_{\mu\nu}$ denotes the Riemannian metric on the embedding spacetime. T is a string tension, a dimension full scale introduced into the theory by hand.

Now instead of using the measure $\sqrt{-\gamma}$, on the 2-dimensional world-sheet, in the framework of this theory two additional worldsheet scalar fields $\varphi^i (i = 1, 2)$ are considered. A new measure density is introduced:

$$\Phi(\varphi) = \frac{1}{2}\epsilon_{ij}\epsilon^{ab}\partial_a\varphi^i\partial_b\varphi^j. \quad (2)$$

There are no limitations on employing any other measure of integration different than $\sqrt{-\gamma}$. The only restriction is that it must be a density under arbitrary diffeomorphisms (reparametrizations) on the underlying spacetime manifold. The modified-measure theory is an example of such a theory.

Then the modified bosonic string action is (as formulated first in [18] and latter discussed and generalized also in [19])

$$S = - \int d^2\sigma \Phi(\varphi) \left(\frac{1}{2}\gamma^{ab}\partial_a X^\mu\partial_b X^\nu g_{\mu\nu} - \frac{\epsilon^{ab}}{2\sqrt{-\gamma}}F_{ab}(A) \right), \quad (3)$$

where F_{ab} is the field-strength of an auxiliary Abelian gauge field A_a : $F_{ab} = \partial_a A_b - \partial_b A_a$.

It is important to notice that the action (3) is invariant under conformal transformations of the internal metric combined with a diffeomorphism of the measure fields,

$$\gamma_{ab} \rightarrow J\gamma_{ab}, \quad (4)$$

$$\varphi^i \rightarrow \varphi'^i = \varphi'^i(\varphi^i) \quad (5)$$

such that

$$\Phi \rightarrow \Phi' = J\Phi \quad (6)$$

Here J is the jacobian of the diffeomorphism in the internal measure fields which can be an arbitrary function of the world sheet space time coordinates, so this can called indeed a local conformal symmetry.

To check that the new action is consistent with the sigma-model one, let us derive the equations of motion of the action (3).

The variation with respect to φ^i leads to the following equations of motion:

$$\epsilon^{ab}\partial_b\varphi^i\partial_a(\gamma^{cd}\partial_c X^\mu\partial_d X^\nu g_{\mu\nu} - \frac{\epsilon^{cd}}{\sqrt{-\gamma}}F_{cd}) = 0. \quad (7)$$

since $\det(\epsilon^{ab}\partial_b\varphi^i) = \Phi$, assuming a non degenerate case ($\Phi \neq 0$), we obtain,

$$\gamma^{cd}\partial_c X^\mu\partial_d X^\nu g_{\mu\nu} - \frac{\epsilon^{cd}}{\sqrt{-\gamma}}F_{cd} = M = \text{const.} \quad (8)$$

The equations of motion with respect to γ^{ab} are

$$T_{ab} = \partial_a X^\mu \partial_b X^\nu g_{\mu\nu} - \frac{1}{2} \gamma_{ab} \frac{\epsilon^{cd}}{\sqrt{-\gamma}} F_{cd} = 0. \quad (9)$$

One can see that these equations are the same as in the sigma-model formulation . Taking the trace of (9) we get that $M = 0$. By solving $\frac{\epsilon^{cd}}{\sqrt{-\gamma}} F_{cd}$ from (8) (with $M = 0$) we obtain the standard string eqs.

The emergence of the string tension is obtained by varying the action with respect to A_a :

$$\epsilon^{ab} \partial_b \left(\frac{\Phi(\varphi)}{\sqrt{-\gamma}} \right) = 0. \quad (10)$$

Then by integrating and comparing it with the standard action it is seen that

$$\frac{\Phi(\varphi)}{\sqrt{-\gamma}} = T. \quad (11)$$

That is how the string tension T is derived as a world sheet constant of integration opposite to the standard equation (1) where the tension is put ad hoc. Let us stress that the modified measure string theory action does not have any *ad hoc* fundamental scale parameters. associated with it. This can be generalized to incorporate super symmetry, see for example [19], [21], [20] , [22]. For other mechanisms for dynamical string tension generation from added string world sheet fields, see for example [27] and [28]. However the fact that this string tension generation is a world sheet effect and not a universal uniform string tension generation effect for all strings has not been sufficiently emphasized before.

Notice that Each String in its own world sheet determines its own tension. Therefore the tension is not universal for all strings.

3 Introducing Background Fields including a New Background Field, The Tension Field

Schwinger [32] had an important insight and understood that all the information concerning a field theory can be studied by understanding how it reacts to sources of different types.

This has been discussed in the text book by Polchinski for example [35] . Then the target space metric and other external fields acquire dynamics which is enforced by the requirement of zero beta functions.

However, in addition to the traditional background fields usually considered in conventional string theory, one may consider as well an additional scalar field that induces currents in the string world sheet and since the current couples to the world sheet gauge fields, this produces a dynamical tension controlled by the external scalar field as shown at the classical level in [33]. In the next two subsections we will study how this comes about in two steps, first we introduce world sheet currents that couple to the internal gauge fields in Strings and Branes and second we define a coupling to an external scalar field by defining a world sheet currents that couple to the internal gauge fields in Strings that is induced by such external scalar field.

3.1 Introducing world sheet currents that couple to the internal gauge fields

If to the action of the string we add a coupling to a world-sheet current j^a , i.e. a term

$$S_{\text{current}} = \int d^2\sigma A_a j^a, \quad (12)$$

then the variation of the total action with respect to A_a gives

$$\epsilon^{ab} \partial_a \left(\frac{\Phi}{\sqrt{-\gamma}} \right) = j^b. \quad (13)$$

We thus see indeed that, in this case, the dynamical character of the brane is crucial here.

3.2 How a world sheet current can naturally be induced by a bulk scalar field, the Tension Field

Suppose that we have an external scalar field $\phi(x^\mu)$ defined in the bulk. From this field we can define the induced conserved world-sheet current

$$j^b = e \partial_\mu \phi \frac{\partial X^\mu}{\partial \sigma^a} \epsilon^{ab} \equiv e \partial_a \phi \epsilon^{ab}, \quad (14)$$

where e is some coupling constant. The interaction of this current with the world sheet gauge field is also invariant under local gauge transformations in the world sheet of the gauge fields $A_a \rightarrow A_a + \partial_a \lambda$.

For this case, (13) can be integrated to obtain

$$T = \frac{\Phi}{\sqrt{-\gamma}} = e\phi + T_i, \quad (15)$$

or equivalently

$$\Phi = \sqrt{-\gamma}(e\phi + T_i), \quad (16)$$

The constant of integration T_i may vary from one string to the other. Notice that the interaction is metric independent since the internal gauge field does not transform under the conformal transformations. This interaction does not therefore spoil the world sheet conformal transformation invariance in the case the field ϕ does not transform under this transformation. One may interpret (16) as the result of integrating out classically (through integration of equations of motion) or quantum mechanically (by functional integration of the internal gauge field, respecting the boundary condition that characterizes the constant of integration T_i for a given string). Then replacing $\Phi = \sqrt{-\gamma}(e\phi + T_i)$ back into the remaining terms in the action gives a correct effective action for each string. Each string is going to be quantized with each one having a different T_i . The consequences of an independent quantization of many strings with different T_i covering the same region of space time will be studied in the next section.

3.3 Consequences from World Sheet Quantum Conformal Invariance on the Tension field, when several strings share the same region of space

3.3.1 The case where all all string tensions are the same, i.e., $T_i = T_0$, and the appearance of a target space conformal invariance

If all $T_i = T_0$, we just redefine our background field so that $e\phi + T_0 \rightarrow e\phi$ and then in the effective action for all the strings the same combination $e\phi g_{\mu\nu}$, and only this combination will be determined by the requirement that the conformal invariance in the world sheet of all strings be preserved quantum mechanically, that is, that the beta function be zero. So in this case we will not be able to determine $e\phi$ and $g_{\mu\nu}$ separately, just the product $e\phi g_{\mu\nu}$, so the equation obtained from equating the beta function to zero will have the target space conformal invariance $e\phi \rightarrow F(x)e\phi$, $g_{\mu\nu} \rightarrow F(x)^{-1}g_{\mu\nu}$.

That is, there is no independent dynamics for the Tension Field in this case. On the other hand, if there are at least two types of string tensions, that symmetry will not exist and there is the possibility of determining separately $e\phi$ and $g_{\mu\nu}$ as we will see in the next subsection.

3.3.2 The case of two different string tensions

If we have a scalar field coupled to a string or a brane in the way described in the sub section above, i.e. through the current induced by the scalar field in the extended object, according to eq. (16), so we have two sources for the variability of the tension when going from one string to the other: one is the integration constant T_i which varies from string to string and the other the local value of the scalar field, which produces also variations of the tension even within the string or brane world sheet.

As we discussed in the previous section, we can incorporate the result of the tension as a function of scalar field ϕ , given as $e\phi + T_i$, for a string with the constant of integration T_i by defining the action that produces the correct equations of motion for such string, adding also other background fields, the anti symmetric two index field $A_{\mu\nu}$ that couples to $\epsilon^{ab}\partial_a X^\mu \partial_b X^\nu$ and the dilaton field φ that couples to the topological density $\sqrt{-\gamma}R$

$$S_i = - \int d^2\sigma (e\phi + T_i) \frac{1}{2} \sqrt{-\gamma} \gamma^{ab} \partial_a X^\mu \partial_b X^\nu g_{\mu\nu} + \int d^2\sigma A_{\mu\nu} \epsilon^{ab} \partial_a X^\mu \partial_b X^\nu + \int d^2\sigma \sqrt{-\gamma} \varphi R. \quad (17)$$

Notice that if we had just one string, or if all strings will have the same constant of integration $T_i = T_0$.

In any case, it is not our purpose here to do a full generic analysis of all possible background metrics, antisymmetric two index tensor field and dilaton fields, instead, we will take cases where the dilaton field is a constant or zero, and the antisymmetric two index tensor field is pure gauge or zero, then the demand of conformal invariance for $D = 26$ becomes the demand that all the metrics

$$g_{\mu\nu}^i = (e\phi + T_i) g_{\mu\nu} \quad (18)$$

will satisfy simultaneously the vacuum Einstein's equations, Notice that if we had just one string, or if all strings will have the same constant of integration $T_i = T_0$, then

all the $g_{\mu\nu}^i$ metrics are the same and then (18) is just a single field redefinition and therefore there will be only one metric that will have to satisfy Einstein's equations, which of course will not impose a constraint on the tension field ϕ .

The interesting case to consider is therefore many strings with different T_i , let us consider the simplest case of two strings, labeled 1 and 2 with $T_1 \neq T_2$, then we will have two Einstein's equations, for $g_{\mu\nu}^1 = (e\phi + T_1)g_{\mu\nu}$ and for $g_{\mu\nu}^2 = (e\phi + T_2)g_{\mu\nu}$,

$$R_{\mu\nu}(g_{\alpha\beta}^1) = 0 \quad (19)$$

and, at the same time,

$$R_{\mu\nu}(g_{\alpha\beta}^2) = 0 \quad (20)$$

These two simultaneous conditions above impose a constraint on the tension field ϕ , because the metrics $g_{\alpha\beta}^1$ and $g_{\alpha\beta}^2$ are conformally related, but Einstein's equations are not conformally invariant, so the condition that Einstein's equations hold for both $g_{\alpha\beta}^1$ and $g_{\alpha\beta}^2$ is highly non trivial.

Let us consider the case that one of the metrics, say $g_{\alpha\beta}^2$ is a Schwarzschild solution, either a 4 D Schwarzschild solution or a product flat of Torus compactified extra dimensions or just a 26 D Schwarzschild solution, in this case, it does not appear possible to have a conformally transformed $g_{\alpha\beta}^2$ for anything else than in the case that the conformal factor that transforms the two metrics is a positive constant, let us call it Ω^2 , in that case $g_{\alpha\beta}^1$ is a Schwarzschild solution of the same type, just with a different mass parameter and different sizes of extra dimensions if the compactified solution is considered. Similar consideration holds for the case the 2 metric is a Kasner solution,

Then in this case also, it does not appear possible to have a conformally transformed $g_{\alpha\beta}^2$ for anything else than in the case that the conformal factor that transforms the two metrics is a constant, we will find other cases where the conformal factor will not be a constant, let us call then conformal factor Ω^2 in general, even when it is not a constant. One can also study metrics used to describe gravitational radiation, then again, multiplying by a constant both the background flat space and the perturbation gives us also a solution of vacuum Einstein's equations.

Then for these situations, we have,

$$e\phi + T_1 = \Omega^2(e\phi + T_2) \quad (21)$$

which leads to a solution for $e\phi$

$$e\phi = \frac{\Omega^2 T_2 - T_1}{1 - \Omega^2} \quad (22)$$

which leads to the tensions of the different strings to be

$$e\phi + T_1 = \frac{\Omega^2(T_2 - T_1)}{1 - \Omega^2} \quad (23)$$

and

$$e\phi + T_2 = \frac{(T_2 - T_1)}{1 - \Omega^2} \quad (24)$$

Both tensions can be taken as positive if $T_2 - T_1$ is positive and Ω^2 is also positive and less than 1. It is important that we were forced to consider a multi metric situation. One must also realize that the constant c is physical, because both metrics live in the

same spacetime, so even if c is a constant, we are not allowed to perform a coordinate transformation, consisting for example of a rescaling of coordinates for one of the metrics and not do the same transformation for the other metric.

Other way to see that Ω^2 is physical consist of considering the scalar consisting of the ratio of the two measures $\sqrt{-g^1}$ and $\sqrt{-g^2}$ where $g^1 = \det(g_{\alpha\beta}^1)$ and $g^2 = \det(g_{\alpha\beta}^2)$, and we find that the scalar $\frac{\sqrt{-g^1}}{\sqrt{-g^2}} = \Omega^D$, showing that Ω is a coordinate invariant.

3.3.3 Flat space in Minkowski coordinates and flat space after a special conformal transformation

Let us study now a case where Ω^2 is not a constant. For this we will consider two spaces related by a conformal transformation, which will be flat space in Minkowski coordinates and flat space after a special conformal transformation.

The flat space in Minkowski coordinates is,

$$ds_1^2 = \eta_{\alpha\beta} dx^\alpha dx^\beta \quad (25)$$

where $\eta_{\alpha\beta}$ is the standard Minkowski metric, with $\eta_{00} = 1$, $\eta_{0i} = 0$ and $\eta_{ij} = -\delta_{ij}$. This is of course a solution of the vacuum Einstein's equations.

We now consider the conformally transformed metric

$$ds_2^2 = \Omega(x)^2 \eta_{\alpha\beta} dx^\alpha dx^\beta \quad (26)$$

which we also demand that will satisfy the D dimensional vacuum Einstein's equations.

Let us use the known transformation law of the Ricci tensor under a conformal transformation applied to $g_{\alpha\beta}^1 = \eta_{\alpha\beta}$ and $\Omega(x)^2 g_{\alpha\beta}^2 = \eta_{\alpha\beta}$, defining $\Omega(x) = \phi^{-1}$, we obtain

$$\begin{aligned} R_{\alpha\beta}^2 = & R_{\alpha\beta}^1 + (D-2)\nabla_\alpha \nabla_\beta (\ln\phi) + \eta_{\alpha\beta} \eta^{\mu\nu} \nabla_\mu \nabla_\nu (\ln\phi) + (D-2)\nabla_\alpha (\ln\phi) \nabla_\beta (\ln\phi) \\ & - (D-2)\eta_{\alpha\beta} \eta^{\mu\nu} \nabla_\mu (\ln\phi) \nabla_\nu (\ln\phi) \end{aligned} \quad (27)$$

Since $g_{\alpha\beta}^1 = \eta_{\alpha\beta}$, we obtain that $R_{\alpha\beta}^1 = 0$, also the covariant derivative above are covariant derivatives with respect to the metric $g_{\alpha\beta}^1 = \eta_{\alpha\beta}$, so they are just ordinary derivatives. Taking this into account, after a bit of algebra we get that,

$$\begin{aligned} R_{\alpha\beta}^2 = & (D-2) \frac{\partial_\alpha \partial_\beta \phi}{\phi} + \eta_{\alpha\beta} \eta^{\mu\nu} \left(\frac{\partial_\mu \partial_\nu \phi}{\phi} - \frac{\partial_\mu \phi \partial_\nu \phi}{\phi^2} \right) \\ & - (D-2) \eta_{\alpha\beta} \eta^{\mu\nu} \frac{\partial_\mu \phi \partial_\nu \phi}{\phi^2} = 0 \end{aligned} \quad (28)$$

by contracting (28) we obtain a relation between $\eta^{\mu\nu} \frac{\partial_\mu \phi \partial_\nu \phi}{\phi}$ and $\eta^{\mu\nu} \partial_\mu \phi \partial_\nu \phi / \phi^2$

$$2\eta^{\mu\nu} \frac{\partial_\mu \partial_\nu \phi}{\phi} = D \frac{\eta^{\mu\nu} \partial_\mu \phi \partial_\nu \phi}{\phi^2} \quad (29)$$

using (29) to eliminate the nonlinear term $\eta^{\mu\nu} \frac{\partial_\mu \phi \partial_\nu \phi}{\phi}$ in (28) we obtain the remarkably simple linear relation,

$$\partial_\alpha \partial_\beta \phi - \frac{1}{D} \eta_{\alpha\beta} \eta^{\mu\nu} \partial_\mu \partial_\nu \phi = 0 \quad (30)$$

So we now first find the most general solution of the linear equation (30), which is,

$$\phi = a_1 + a_2 K_\mu x^\mu + a_3 x^\mu x_\mu \quad (31)$$

and then impose the non linear constraint (29), which implies,

$$a_1 = \frac{a_2^2 K_\mu K^\mu}{4a_3} \quad (32)$$

we further demand that $\phi(x^\mu = 0) = 1$, so that,

$$\phi = 1 + a_2 K_\mu x^\mu + \frac{a_2^2 K_\mu K^\mu}{4} x^\mu x_\mu \quad (33)$$

This coincides with the results of Culetu [36] for $D = 4$ and to identify this result with the result of a special conformal transformation, see discussions in [37] and [38], to connect to standard notation we identify $a_2 K_\mu = 2a_\mu$, so that

$$\phi = 1 + 2a_\mu x^\mu + a^2 x^2 \quad (34)$$

where $a^2 = a^\mu a_\mu$ and $x^2 = x^\mu x_\mu$.

In this case, this conformal factor coincides with that obtained from the special conformal transformation

$$x'^\mu = \frac{(x^\mu + a^\mu x^2)}{(1 + 2a_\nu x^\nu + a^2 x^2)} \quad (35)$$

As discussed by Zumino [38] the finite special conformal transformation mixes up in a complicated way the topology of space time, so it is not useful to interpret the finite special conformal transformations as mapping of spacetimes.

In summary, we have two solutions for the Einstein's equations, $g_{\alpha\beta}^1 = \eta_{\alpha\beta}$ and

$$g_{\alpha\beta}^2 = \Omega^2 \eta_{\alpha\beta} = \phi^{-2} \eta_{\alpha\beta} = \frac{1}{(1 + 2a_\mu x^\mu + a^2 x^2)^2} \eta_{\alpha\beta} \quad (36)$$

We can then study the evolution of the tensions using $\Omega^2 = \phi^{-2} = \frac{1}{(1 + 2a_\mu x^\mu + a^2 x^2)^2}$. We will consider two different cases: 1) $a^2 = 0$, 2) $a^2 \neq 0$

3.3.4 Light Like Segment Compactification

Here we consider the case $a^2 = 0$, and let us consider $a^\mu = (A, A, 0, \dots, 0)$. Then

$$\Omega^2 = \frac{1}{(1 + 2a_\mu x^\mu)^2} = \frac{1}{(1 + 2A(t - x))^2} \quad (37)$$

From this, let us calculate the tensions of the two string types and see that they will be constrained to be inside a segment that moves with the speed of light. At the boundaries of those segments the string tensions become infinity, so the strings cannot escape this segment.

37 leads to the tensions of the different strings to be

$$e\phi + T_1 = \frac{\Omega^2(T_2 - T_1)}{1 - \Omega^2} = \frac{(T_2 - T_1)(1 + 2A(t - x))^2}{4A(t - x)(1 + A(t - x))} \quad (38)$$

and

$$e\phi + T_2 = \frac{(T_2 - T_1)}{1 - \Omega^2} = \frac{(T_2 - T_1)}{4A(t-x)(1 + A(t-x))} \quad (39)$$

Let us take $T_2 - T_1$ positive, A negative, so we see that both tensions above go to positive infinity when $t - x$ goes to zero from negative values. Also both tensions above go to positive infinity when $t - x$ goes to the value $-1/A$ from above. That means that the strings are confined to the moving segment where $t - x$ is inside the segment $(-1/A, 0)$. We call this phenomenon "Light Like Segment Compactification". Light like extra dimensions have been considered by Searight [39] and Braneworlds via Lightlike Branes was considered in [40] for example.

In the next subsection we will see how thick, expanding at sub luminal velocities braneworld scenarios are obtained from Dynamical String Tension Theories.

3.3.5 Braneworlds in Dynamical String Tension Theories

We now consider the case when a^μ is not light like and we will find that for $a^2 \neq 0$, irrespective of sign, i.e. irrespective of whether a^μ is space like or time like, we will have thick Braneworlds where strings can be constrained between two concentric spherically symmetric bouncing higher dimensional spheres and where the distance between these two concentric spherically symmetric bouncing higher dimensional spheres approaches zero at large times. The string tensions of the strings one and two are given by

$$e\phi + T_1 = \frac{(T_2 - T_1)(1 + 2a_\mu x^\mu + a^2 x^2)^2}{(1 + 2a_\mu x^\mu + a^2 x^2)^2 - 1} = \frac{(T_2 - T_1)(1 + 2a_\mu x^\mu + a^2 x^2)^2}{(2a_\mu x^\mu + a^2 x^2)(2 + 2a_\mu x^\mu + a^2 x^2)} \quad (40)$$

$$e\phi + T_2 = \frac{(T_2 - T_1)}{(1 + 2a_\mu x^\mu + a^2 x^2)^2 - 1} = \frac{(T_2 - T_1)}{(2a_\mu x^\mu + a^2 x^2)(2 + 2a_\mu x^\mu + a^2 x^2)} \quad (41)$$

Then, the locations where the string tensions go to infinity are determined by the conditions

$$2a_\mu x^\mu + a^2 x^2 = 0 \quad (42)$$

or

$$2 + 2a_\mu x^\mu + a^2 x^2 = 0 \quad (43)$$

Let us start by considering the case where a^μ is time like, then without losing generality we can take $a^\mu = (A, 0, 0, \dots, 0)$. In this case the denominator in (40), (41) is

$$(2a_\mu x^\mu + a^2 x^2)(2 + 2a_\mu x^\mu + a^2 x^2) = (2At + A^2(t^2 - x^2))(2 + 2At + A^2(t^2 - x^2)) \quad (44)$$

The condition (42) implies then that

$$x_1^2 + x_2^2 + x_3^2 \dots + x_{D-1}^2 - (t + \frac{1}{A})^2 = -\frac{1}{A^2} \quad (45)$$

while the other boundary of infinite string tension (43) is given by,

$$x_1^2 + x_2^2 + x_3^2 \dots + x_{D-1}^2 - (t + \frac{1}{A})^2 = \frac{1}{A^2} \quad (46)$$

So we see that (49) represents an exterior boundary which has an bouncing motion with a minimum radius $\frac{1}{A}$ at $t = -\frac{1}{A}$, The denominator (??) is positive between

these two bubbles. So for $T_2 - T_1$ positive the tensions are positive and diverge at the boundaries defined above.

The internal boundary (45) exists only for times t smaller than $-\frac{2}{A}$ and bigger than 0, so in the time interval $(-\frac{2}{A}, 0)$ there is no inner surface of infinite tension strings. This inner surface collapses to zero radius at $t = -\frac{2}{A}$ and emerges again from zero radius at $t = 0$.

For large positive or negative times, the difference between the upper radius and the lower radius goes to zero as $t \rightarrow \infty$

$$\sqrt{\frac{1}{A^2} + (t + \frac{1}{A})^2} - \sqrt{-\frac{1}{A^2} + (t + \frac{1}{A})^2} \rightarrow \frac{1}{tA^2} \rightarrow 0 \quad (47)$$

of course the same holds $t \rightarrow -\infty$. This means that for very large early or late times the segment where the strings would be confined (since they will avoid having infinite tension) will be very narrow and the resulting scenario will be that of a brane world for late or early times, while in the bouncing region the inner surface does not exist. Let us start by considering the case where a^μ is space like, then without losing generality we can take $a^\mu = (0, A, 0, \dots, 0)$. In this case the denominator in (40), (41) is

$$(2a_\mu x^\mu + a^2 x^2)(2 + 2a_\mu x^\mu + a^2 x^2) = (-2Ax^1 - A^2(t^2 - \vec{x}^2))((2 - 2Ax^1 - A^2(t^2 - \vec{x}^2))) \quad (48)$$

where $\vec{x} = (x^1, x^2, \dots, x^{D-1})$ represents the spacial part of x^μ , and $\vec{x}^2 = (x^1)^2 + (x^2)^2 + \dots + (x^{D-1})^2$. We now consider the case when a^μ is space like, then without losing generality we can take $a^\mu = (0, A, 0, \dots, 0)$. We then consider the first boundary where the string tensions approach infinity according to (42),

$$-(x_1 - \frac{1}{A})^2 - x_2^2 - x_3^2 \dots - x_{D-1}^2 + t^2 = -\frac{1}{A^2} \quad (49)$$

which describes a bouncing bubble with minimum radius $\frac{1}{A}$ at $t = 0$.

The case (43) gives

$$-(x_1 - \frac{1}{A})^2 - x_2^2 - x_3^2 \dots - x_{D-1}^2 + t^2 = \frac{1}{A^2} \quad (50)$$

(50) is an internal boundary which exists only for times t smaller than $-\frac{1}{A}$ and bigger than $\frac{1}{A}$. Between $-\frac{1}{A}$ and $\frac{1}{A}$ there is no inner surface of infinite tension strings. Between these two bubbles the two factors in eq. This inner surface collapses to zero radius at $t = -\frac{1}{A}$ and emerges again from zero radius at $t = \frac{1}{A}$. So the situation is very similar to that of the case where the vector a^μ is time like, just that the roles of the cases $\Omega = 1$ and $\Omega = -1$ get exchanged. Between these two boundaries the two factors in the denominator (48) are positive, while at the boundaries one or the other approach zero and the tensions diverge, so again for $T_2 - T_1$ positive the tensions are positive and diverge at the boundaries.

Once again for large positive or negative times, the difference between the upper radius and the lower radius goes to zero. Implying that the strings will be confined to a very small segment at large early or late times, so then again we get an emergent brane world scenario.

The strings and therefore all matter and gravity will be consequently confined to the very small segment of size $\frac{1}{tA^2}$, very small for large t . At the moment of the bounce there is no brane world, there is only one exterior bubble which represents infinite

tension location, the brane is generated dynamically after a period of time by the appearance of the inner bubble which completes the trapping of the strings between two surfaces.

Our approach is very different to the present standard approaches to braneworlds in the context of string theories however. In our approach a dynamical string tension has been used. Our scenario could be enriched by incorporating aspects of the more traditional braneworlds, like introducing D-branes between the surfaces where the string tensions go to infinity, so open strings could end before their tensions approach an infinite value, or the surfaces where the tensions diverge could be themselves be defined as D branes for open strings. These possibilities have been ignored here to simplify the discussion.

In any case, given that the tension of the strings diverge at the two boundaries we have defined, all strings are confined between those, the closed strings also, so unlike more traditional braneworlds, gravity does not escape to the bulk, in fact in the framework proposed here a braneworld scenario using just closed strings is perfectly possible.

Finally, since the string tensions go to infinity at the boundaries, there is the option of avoiding the Hagedorn Temperature, which is proportional to the string tension, as it was discussed in the case of other examples where the string tensions go to infinity [26].

Acknowledgments I thank Oleg Andreev, David Andriot, Stefano Ansoldi, David Benisty, Euro Spallucci, Emil Nissimov, Svetlana Pacheva, Tatiana Vulfs, Hitoshi Nishino, Subhash Rajpoot, Luciano Rezzolla, Horst Stöcker, Jürgen Struckmeier, David Vasak, Johannes Kirsch, Dirk Kehm and Matthias Hanauske for useful discussions. I also want to thank the Foundational Questions Institute (FQXi) and the COST actions Quantum Gravity Phenomenology in the multi messenger approach, CA18108 and Gravitational waves, Black Holes and Fundamental Physics, CA16104 for support and special thanks to the Frankfurt Institute for Advanced Study for hospitality and financial support and to Astrophysics Group at Goethe University for providing me with the opportunity of presenting these results in a seminar at one of their Astro Coffe Seminars, <https://astro.uni-frankfurt.de/astrocoffee/>.

References

- [1] Rubakov, V. A.; Shaposhnikov, M. E. (1983). "Do we live inside a domain wall?". *Physics Letters. B.* 125 (2–3): 136–138.
- [2] M Visser, An exotic class of Kaluza-Klein models, *Letters B*, 1985
- [3] Randall, Lisa; Sundrum, Raman (1999). "Large Mass Hierarchy from a Small Extra Dimension". *Physical Review Letters.* 83 (17): 3370–3373. arXiv:hep-ph/9905221.
- [4] Randall, Lisa; Sundrum, Raman (1999). "An Alternative to Compactification". *Physical Review Letters.* 83 (23): 4690–4693. arXiv:hep-th/9906064
- [5] Matej Pavsic, A Brane world model with intersecting branes, *Phys.Lett.A* 283 (2001) 8 • e-Print: hep-th/0006184 [hep-th]
- [6] Gogberashvili, M. "Hierarchy problem in the shell universe model". *International Journal of Modern Physics D.* 11 (10): 1635–1638. arXiv:hep-ph/9812296

- [7] Intersecting Brane Worlds - A Path to the Standard Model? by Dieter Lust (hep-th/0401156, 31 pages)
- [8] "Superstrings", John H. Schwarz, Vols 1 and 2, World Scientific, 1985; M. B. Green, J. H. Schwarz and E. Witten, Superstring Theory, Cambridge University Press, 1987.
- [9] E.I. Guendelman, A.B. Kaganovich, Phys.Rev.D55:5970-5980 (1997)
- [10] E.I. Guendelman, Mod.Phys.Lett.A14, 1043-1052 (1999)
- [11] E.I. Guendelman, O. Katz, Class.Quant.Grav. 20 (2003) 1715-1728 • e-Print: gr-qc/0211095 [gr-qc]
- [12] Frank Gronwald, Uwe Muench, Alfredo Macias, Friedrich W. Hehl, Phys.Rev.D 58 (1998) 084021 • e-Print: gr-qc/9712063 [gr-qc]
- [13] Eduardo Guendelman, Ramón Herrera, Pedro Labrana, Emil Nissimov, Svetlana Pacheva, Gen.Rel.Grav. 47 (2015) 2, 10 • e-Print: 1408.5344 [gr-qc]
- [14] Eduardo Guendelman, Douglas Singleton, Nattapong Yongram, JCAP 11 (2012) 044 • e-Print: 1205.1056 [gr-qc]
- [15] R. Cordero, O.G. Miranda, M. Serrano-Crivelli, JCAP 07 (2019) 027 • e-Print: 1905.07352 [gr-qc]
- [16] Eduardo Guendelman, Emil Nissimov, Svetlana Pacheva, Eur.Phys.J.C 75 (2015) 10, 472 • e-Print: 1508.02008 [gr-qc]
- [17] Conformally invariant gauge theory of three-branes in 6-D and the cosmological constant, E.I. Guendelman, E. Spallucci, Phys.Rev.D 70 (2004) 026003 • e-Print: hep-th/0311102 [hep-th], Conformally invariant brane world and the cosmological constant, E.I. Guendelman, Phys.Lett.B 580 (2004) 87-92 • e-Print: gr-qc/0303048 [gr-qc]
- [18] E.I. Guendelman, Class.Quant.Grav. 17, 3673-3680 (2000)
- [19] E.I. Guendelman, A.B. Kaganovich, E.Nissimov, S. Pacheva, Phys.Rev.D66:046003 (2002)
- [20] E.I. Guendelman, Phys.Rev.D 63 (2001) 046006 • e-Print: hep-th/0006079 [hep-th]
- [21] Hitoshi Nishino, Subhash Rajpoot, Phys.Lett.B 736 (2014) 350-355 e-Print: 1411.3805 [hep-th].
- [22] T.O. Vulfs, E.I. Guendelman, Annals Phys. 398 (2018) 138-145 • e-Print: 1709.01326 [hep-th]
- [23] T.O. Vulfs, E.I. Guendelman, Int.J.Mod.Phys.A 34 (2019) 31, 1950204 • e-Print: 1802.06431 [hep-th]
- [24] T.O. Vulfs, Ben Gurion University Ph.D Thesis, (2021), arXiv:2103.08979.
- [25] E.I. Guendelman, Cosmology and Warped Space Times in Dynamical String Tension Theories, e-Print: 2104.08875
- [26] E.I. Guendelman, Escaping the Hagedorn Temperature in Cosmology and Warped Spaces with Dynamical Tension Strings, e-Print: 2105.02279 [hep-th].
- [27] P.K. Townsend, Phys.Lett.B 277 (1992) 285-288.

- [28] E. Bergshoeff, L.A.J. London, P.K. Townsend, *Class.Quant.Grav.* 9 (1992) 2545-2556, *Class. Quantum Grav.* 9 (1992) 2545-2556 • e-Print: hep-th/9206026 [hep-th]
- [29] Deser, S. and Zumino, *Phys. Lett.* B65 , 369, (1976)
- [30] Brink , L., Di Vecchia, P and Howe, S. , *Phys. Lett.* B65, 471 , (1976)
- [31] Polyakov, A. M., , *Phys. Lett.* B103 ,207, (1980).
- [32] *Particles and Sources*, Julian Schwinger, *Phys.Rev.* 152 (1966) 1219-1226 DOI: 10.1103/PhysRev.152.1219
- [33] S. Ansoldi, E. I. Guendelman, E. Spallucci, *Mod.Phys.Lett.A* 21 (2006) 2055-2065 • e-Print: hep-th/0510200 [hep-th]
- [34] A. Vilenkin, E.P. S. Shellard, *Cosmic Strings and Other Topological Defects*, Cambridge Monographs on Mathematical Physics, Cambridge University Press, 2000, 580 pages, ISBN: 9780521654760.
- [35] Joseph Polchinski, *String Theory*, vol. 1 , Cambridge University Press (1998) ; some papers on strings with background fields are C. G. Callan, D. Friedan, E. J. Martinec and M. J. Perry, *Nucl. Phys. B* 262 (1985) 593; T. Banks, D. Nemeschansky and A. Sen, *Nucl. Phys. B* 277 (1986) 67
- [36] H. Culetu, The special conformal transformation and Einstein's equations, *Il Nuovo Cimento B*, 621–628 (1989)
- [37] On the Advancements of Conformal Transformations and their Associated Symmetries in Geometry and Theoretical Physics1, H.A. Kastrup, *Annalen Phys.*17:631-690,2008, arXiv:0808.2730 [physics.hist-ph].
- [38] *Effective Lagrangians and Broken Symmetries* B. Zumino (1970), in: *Lectures on Elementary Particles and Quantum Field Theory v.2*, Cambridge, Mass.: Brandeis Univ., pp. 437-500
- [39] On degenerate metrics and electromagnetism T.P. Searight, *Gen.Rel.Grav.* 35 (2003) 791-805 • e-Print: hep-th/0405204 [hep-th]
- [40] *Space-Time Compactification, Non-Singular Black Holes, Wormholes and Braneworlds via Lightlike Branes*, Eduardo Guendelman, Alexander Kaganovich, Emil Nissimov, Svetlana Pacheva, in: , "Sixth Mathematical Physics Meeting", pp. 217-234, B. Dragovic and Z. Rakic (eds.), Belgrade Inst. Phys. Press (2011) • Contribution to: MPHYS6, 217-233 • e-Print: 1011.6241 [hep-th]